

AD AO 62838

DDC FILE COPY

LEVEL

OFFICE OF NAVAL RESEARCH

Contract No. Nggo14-76-C-9399

Task No. NR 053-608

TECHNICAL REPORT NO. 103

(10

6 Denterin rept

Characterization and Molecular Structure of [(PPh'3)RhC2B9H11]2,

a Phosphinorhodacarborane Dimer Containing Rn-H-B Bridges •

)

R. T. Baker, R. E. King, C. Knobler, C. A. O'Con M. F. Hawthorne

Prepared for Publication

ir

Journal of the American Chemical Society

29 Nov 78

Department of Chemistry
University of California
Los Angeles, California 90024


DDC DEC 29 1978 F

November 20, 1978

Reproduction in whole or part is permitted for any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

78 12 27 0067B



Characterization and Molecular Structure of [(PPh3)RhC2B9H11]2,

a Phosphinorhodacarborane Dimer Containing Rh-H-B Bridges

by

R. T. Baker, R. E. King, C. Knobler, C. A. O'Con and M. F. Hawthorne\*

Department of Chemistry University of California Los Angeles, California 90024 We have recently shown that a number of B-H containing substrates undergo catalytic isotopic exchange of terminal hydrogen with deuterium gas in the presence of the hydridorhodacarborane [3,3-(PPh<sub>3</sub>)<sub>2</sub>-3-H-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] (I) and a mechanism was postulated which involved the oxidative addition of boron-hydrogen bonds to the catalytic species. The complex I is also an efficient homogeneous catalyst for the hydrogenation of terminal olefins and it was observed during hydrogenation studies that exposure of a yellow solution of I and substrate olefin to air produced a purple product. We wish to report the complete characterication of the purple complex (II), which can be viewed as a model intermediate in the oxidative addition of a terminal boron-hydrogen bond to rhodium.

The complex(II) was prepared by heating a solution of (I) in benzene with 2 equivalents of benzoyl peroxide. The reaction mixture was purified by column chromatography (silica gel/2:1/heptane-benzene) and the complex obtained in 40% yield as dark purple microcrystals.

The infrared spectrum of II (nujol) showed a band characteristic of terminal B-H bonds at 2600 cm<sup>-1</sup> and peaks due to the coordinated dicarbollide and triphenylphosphine ligands. The visible spectrum in benzene exhibited peaks at 570 nm, (log  $\varepsilon$  = 3.76) and 390 nm(3.76) and the elemental analysis supported the formulation of (II) as  $[(Ph_3P)RhC_2B_9H_{11}]_n$  which resembled a partially characterized compound previously reported in the literature. [Anal. Found: C, 48.27; H, 5.21; B, 19.76; P, 5.72; Rh, 20.01; Calcd for  $C_{20}H_{26}PRhB_9$ : C, 48.28; H, 5.27; B, 19.55; P, 6.22; Rh, 20.68]. Osmometric molecular weight studies in benzene gave MW = 965  $\pm$  32 (3 determinations). Calcd for  $[C_{20}H_{26}PRhB_9]_2$ : 995.22. The 200 MHz  $^1$ H NMR spectrum in  $CD_2C1_2$  showed a complex multiplet centered at 7.336 due to the PPh3 ligands, three carborane C-H resonances at 3.52 (area 2), 2.57(1) and 1.62(1)6 and a complex multiplet at 1.236 due to B-H resonances. A broad resonance at -18.56 due to the hydrogen atoms of the inequivalent Rh-H-B bridges

sharpens to two resonances at 233°K (-18.0 and -19.06). The 80.5 MHz  $^{11}$ B{ $^{1}$ H} NMR spectrum in CD<sub>2</sub>Cl<sub>2</sub>  $^{9}$  exhibited peaks at -26.2, -16.3, -4.8, 3.5 and 24.2 ppm in a ratio of 1:8:7:1:1. The 81.02 MHz  $^{31}$ P{ $^{1}$ H} NMR spectrum in C<sub>6</sub>D $^{10}_{6}$  gave two doublets of equal intensity centered at 36.7 ppm, J<sub>P-Rh</sub>=172 Hz;  $^{11}$  and 29.7 ppm, J<sub>P-Rh</sub>=135 Hz.

Suitable crystals for x-ray diffraction studies were obtained by layering a  $\text{CH}_2\text{Cl}_2$  solution of II with heptane. The dark purple crystals of II are triclinic,  $\overline{\text{Pl}}$ . Cell data are presented in Table 1. The molecular structure of

#### Table I

II is shown in Figure 1; significant bond distances and angles are presented in Table 2.

Figure 1

Table II

As can be seen from Figure 1, each rhodium atom is symmetrically bound to the pentagonal face of the dicarbollide ligand and interacts with the other dicarbollide ligand via a three-center Rh-H-B bridge. The Rh-H distances of 1.77(6) and 1.78(6) Å compare well with those in other M-H-B bridges such as the Co-H distance of 1.82(7) Å found in  $[(C_2H_5)_4N][Co[(C_2B_10H_{10})_2]_2]^{15}$  and the Cr-H distance of 1.78(6) Å found in  $[(CH_3)_4N][(CO)_4CrB_3H_8]$ . A Rh-H distance of 2.1 Å has been reported for the Rh-H-B bridge in  $[(PPh_3)_2Rh(CB_{10}H_{10}C-C_6H_5)]$ . The Rh-Rh distance of 2.763(1) is well within the range reported for Rh-Rh single bonds (cf.  $Rh^I$ - $Rh^I$  = 2.68 Å in  $[(n^5-C_5H_5)_2Rh_2(CO)_3]$ ,  $Rh^{III}$ - $Rh^{III}$  = 2.906(1) in  $[(\mu$ -H) $(\mu$ -C1) $\{(n^5-C_5(CH_3)_5)RhC1\}_2]$ ).

The transformation of a terminal B-H bond in I into a Rh-H-B bridge in II may represent an intermediate step in the oxidative addition of terminal B-H bonds to Rh. A similar situation is seen in the molecular structure of  $[Rh(PPh_3)_3](Clo_4)$ , on which the metal atom appears to interact with the C-H bond of a phenyl ring  $(Rh-H=2.56\ \text{Å})$ . The two terminal B-H bonds involved in the Rh-H-B interactions determine a specific stereoisomer and of the four possibilities only II is observed with both d- and l-enantiomers present in the unit cell.

Homogeneous olefin hydrogenation experiments were performed to determine the catalytic activity of II. <sup>22</sup> In typical experiments it was found that II is an active hydrogenation catalyst exhibiting rates comparable to I, per rhodium center. <sup>23</sup> Further studies concerning the mechanism of homogeneous hydrogenation of olefin utilizing I and II as catalysts are presently underway in these laboratories.

### Acknowledgements

The authors wish to thank Professor F. A. L. Anet for the use of the 80.5 MHz <sup>11</sup>B NMR spectrometer, Ms. Betsy Anfield for the preliminary kinetic data, Drs. J. J. Wilczynski and S. B. Miller for their preliminary work on II and Mr. B. A. Katz for assistance with the crystal structure. We also wish to thank the NSF for the purchase of the Bruker WP-200 NMR Spectrometer (Grant No. CHE76-05926) and the Syntex Diffractometer (Grant No. GP28248). This research was supported in part by the Office of Naval Research.

TABLE I

Cell Data		298 <sup>0</sup> K	113 <sup>0</sup> K
Cell Dimensions	<u>a</u> -	11.202(2) A	11.118(2)
	<u>b</u> =	13.660(3)	13.456(3)
	<u>c</u> =	18.565(3)	18.390(3)
	α <b>=</b>	93.48(2) <sup>0</sup>	93.09(2)
	β=	76.44(1)	76.22(1)
	Υ=	76.80(2)	76.90(2)
Cell Volume	V=	2669.1(1.0) Å <sup>3</sup>	2585.7(9)
X-ray density	ρ=	1.350 gmcm <sup>-3</sup> ; Z=2	1.362
Flotation density	KI (aq)	1.32(5)gmcm <sup>-3</sup>	

TABLE II

	Se	elected Bond	Distances and	Angles	
Distances	(Å) - Rh(3)	Rh(3')	Angles (°)	Rh(3)	Rh(3')
Rh-H	1.78(6)	1.77(6)	Rh-H-B <sub>g</sub>	97(4)	102(5)
Rh-B <sub>g</sub>	2.327(8)	2.238(8)	Rh-Bg-Rh*	75.7(2)	77.2(2)
Rh-C	2.251(6) 2.280(7)	2.214(7) 2.269(7)	B <sub>o</sub> -Rh-P	101.1(2)	96.2(2)
Rh-B	(2.190(8) {2.211(8)	2.173(8) 2.190(8)	Rh-Rh*-B <sub>o</sub>	50.6(2)	49.7(2)
Rh-P	(2.220(8) 2.338	2.190(8) 2.221(8) 2.360(2)	P-Rh-H	85(2)	89(2)
Rh-Rh	2.7628	(9)			


<sup>\*</sup>Refers to Rh atom not indicated by column heading.

#### REFERENCES

- E. L. Hoel, M. Talebinasab-Savari, and M. F. Hawthorne, <u>J. Am. Chem. Soc.</u>, 99, 4356 (1977).
- 2. T. E. Paxson and M. F. Hawthorne, J. Am. Chem. Soc., 96, 4674 (1974).
- 3. Mechanistic hydrogenation studies are presently underway utilizing an automatic gas titration apparatus to be described later. B. A. Anfield, M. S. Delaney, and M. F. Hawthorne (to be published).
- 4. This color change is also effected by dissolution of I in wet reagent grade acetonitrile or tetrahydrofuran and is accelerated in benzene by heating with benzoyl peroxide.
- 5.  $[closo 3, 3' (PPh_3)_2 3, 8'; 3', 7 \mu H, H' 3, 8'; 3', 7 Rh_2 1, 2; 1', 2' (C_2B_9H_{11})_2]$ .
- 6. The compound is soluble in most organic solvents except alcohols and alkanes; R<sub>f</sub>=0.65 (benzene/silica gel). [Melting point: 187-189<sup>0</sup>C, <u>in vacuo</u> (uncorrected)].
- 7. A. R. Siedle, <u>J. Organomet. Chem.</u>, 249 (1975). We have repeated this work and shown that the " $[(C_2B_9H_{11})Rh(PPh_3)C_6H_6]_2$ " compound is in fact the benzene solvate of II,  $[(PPh_3)Rh(C_2B_9H_{11})]_2 \cdot 2C_6H_6$ , and the reported visible spectrum is in error (<u>ca</u>. 50 nm). Crystals of the solvate suitable for x-ray work could not be obtained due to loss of benzene which rapidly destroys the crystal.
- 8. Examination of Figure 1 indicates that while C(2) and C(1) are  $\alpha$  and  $\beta$  (respectively) to the boron atom (B(7)) involved in the Rh-H-B bridge, C(2') and C(1') are both  $\beta$  to B(8') and thus the  $^1$ H NMR chemical shifts of H(2') and H(1') are indistinguishable at 200 MHz and a resonance of area 2 is observed.
- 9. Referenced to the external standard Et<sub>2</sub>0·BF<sub>3</sub>.
- 10. Referenced to the external standard D3PO4.
- 11. A further splitting of 5Hz is observed for this resonance due to <sup>31</sup>P coupling to the Rh-H-B hydrogen nucleus. This splitting is removed by suitably offsetting the broad-band <sup>1</sup>H decoupling frequency.
- 12. The crystals contain two molecules of CH<sub>2</sub>Cl<sub>2</sub> per molecule of II and were coated with epoxy resin for x-ray work.

- 13. X-ray intensity data were collected by the 0-20 technique with MoK $_{\alpha}$  radiation [graphite monochromator] on a Syntex PI automated diffractometer equipped with a scintillation counter and a pulse height analyzer. The low temperature attachment (ref. 14) maintained the crystal temperature at -160  $\pm$  1°C with a temperature stability of  $\pm$  0.5°C. Of a total of 6388 reflections examined, 5535 had I > 3 $\sigma$ (I) and were used in the structure determination. The data were corrected for absorption, Lorentz and polarization effects. The structure was solved by heavy atom methods and refined by anisotropic least-squares refinement, converging at R = 0.047 and R = 0.072.
- 14. C. E. Strouse, Rev. Sci. Inst., 47, 871 (1976).
- 15. R. A. Love and R. Bau, J. Am. Chem. Soc., 94, 8274 (1972).
- 16. L. J. Guggenberger, Inorg. Chem., 9, 367 (1970).
- 17. G. Allegra, M. Calligaris, R. Furlanetto, G. Nardin and L. Randaccio, <u>Cryst.</u>
  <u>Struct. Comm.</u>, 3, 69 (1974).
- 18. O. S. Mills and J. P. Nice, J. Organomet. Chem., 10, 337 (1967).
- 19. M. R. Churchill and S. W. Y. Ni, J. Am. Chem. Soc., 95, 2150 (1973).
- 20. Y. W. Yared, S. L. Miles, R. Bau, and C. A. Reed, ibid., 99, 7076 (1977).
- 21. Current investigation of mixed aryl-alkyl phosphine derivatives of I indicates that an identical stereoselective decomposition occurs to give analogs of II. R. T. Baker and M. F. Hawthorne (to be published).
- 22. Potential catalysts are screened in purified tetrahydrofuran solvent utilizing a gas-tight glass apparatus equipped with an oil diffusion pump, 100 ml Schlenk flask, H<sub>2</sub> inlet, mercury manometer and bubbler. The olefin is injected through a septum on the side-arm using a Gilmont ultra precision syringe and H<sub>2</sub> (1 atm initial pressure) uptake is recorded as a function of time. The reaction solution must be rapidly stirred to avoid diffusion control of the hydrogenation rates.
- 23. Using an initial hydrogen pressure of 1 atm, an initial vinyltrimethylsilane concentration of 9.84 x  $10^{-3}$ M and catalyst concentrations of 2.72 x  $10^{-3}$ M, the initial hydrogenation rates ( $\pm$  10%) at 24.6°C for I and II were 5.36 x  $10^{-5}$  and 1.27 x  $10^{-4}$  mol/sec, respectively.

Figure 1 Molecular Structure of II. (CH<sub>2</sub>Cl<sub>2</sub> solvate molecules and phenyl rings on phosphorus atoms are omitted for clarity)



UNCLASSIFIED

REPORT DOCUMENTATION	READ INSTRUCTIONS BEFORE COMPLETING FORM	
Technical Report 103	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
Characterization and Molecular Structure of [(PPh3)RhC2B9H11]2, a Phosphinorhodacarborane		S. TYPE OF REPORT & PERIOD COVERED  Interim  6. PERFORMING ORG. REPORT NUMBER
Dimer Containing Rh-H-B Bridges		
7. AUTHOR(+)		S. CONTRACT OR GRANT NUMBER(4)
R. T. Baker, R. E. King, C. Knobl and M. F. Hawthorne	er, C. A. O'Con	N00014-76-C-0390
9. PERFORMING ORGANIZATION HAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
The University of California Department of Chemistry Los Angeles, California 90024		NR 053-608
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
Chemistry Branch		November 20, 1978
Office of Naval Research		13. NUMBER OF PAGES
Washington, D.C. 20360		7 (incl 1 fig.) 15. SECURITY CLASS. (of this report)
14. MONITORING AGENCY NAME & ADDRESS(II differen	t from Controlling Office)	
		Unclassified
		SCHEOULE
16. DISTRIBUTION STATEMENT (of this Report)		

Approved for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Dimeric rhodacarborane Hydrogenation catalyst

The decomposition of Rh(H)(PPh3)2C2B9H1(I) yields [Rh(PPh3)C2B9H11]2 (II) as the only tractable product. The spectral characterization and x-ray molecular structure determination indicate that a terminal B-H bond in (I) has become a Rh-H-B bridge in (II) and (II) is thus a model intermediate in the oxidative addition of a terminal B-H bond to rhodium, the catalytic activity of (II) for the homogeneous hydrogenation of olefins is comparable to that of (I) per rhodium center.

APPENDIX

## TECHNICAL REPORT DISTRIBUTION LIST

No.	Copies	No.	Copies
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 . Attn: Dr. Jerry Smith	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office 760 Market Street, Rm. 447 San Francisco, California 9410 Attn: Dr. P. A. Miller	2 1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller Dr. R. M. Grimes	1
ONR Branch Office 666 Summer St., Bildg. 114, Sec. Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	D 1	University of Virginia Department of Chemistry Charlottesville, Virginia 22901 Dr. M. Tsutsui	1
Director, Naval Research Labora Washington, D.C. 20390 Attn: Code 6100	tory 1	Texas A&M University Department of Chemistry College Station, Texas 77843 Dr. C. Quicksall	1
The Asst. Secretary of the Navy Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	(R&D)	Georgetown University Department of Chemistry 37th & O Streets Washington, D.C. 20007	1
Commander, Naval Air Systems Commander, Naval Air Systems Commander, Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasse)		Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130	
Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1	Washington, D.C. 20375	1

# No. Copies

Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1
Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1
Dr. M. H. Chisholm Princeton University Department of Chemistry Princeton, New Jersey 08540	1
Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1 —
Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1
Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code Washington, D.C. 20380	RD-1)
Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1